



# Binge Drinking Among College Students & Young Adults

## Know the Facts: Binge Drinking

### What is binge drinking?



4 or more drinks (female)



5 or more drinks (male)

on one occasion, generally within about two hours\*

### A standard drink is:



OR



OR



1.5 ounces of distilled spirits or liquor

12 ounces of beer

5 ounces of wine

1.5 ounces of distilled spirits or liquor

\*The above definition is given for a typical adult but may vary depending upon an individual's body weight

### Wisconsin Statistics

47%

of 18-25 year-olds in Wisconsin reported binge drinking at least once within the past month

35%

of all University of Wisconsin (UW) system college students reported binge drinking at least once within the past 2 weeks

23%

of UW system college students reported driving after drinking any alcohol within the past month\*

35%

of UW system college students who drank alcohol reported doing something they later regretted when drinking during the past year

Source: NCHA, 2015 & NSDUH, 2016-17

\*Students responding "N/A, don't drive" and "N/A, don't drink" were excluded from the analysis

## The Dangers of Spring Break & Binge Drinking

College students are at **increased risk** for binge drinking, which can lead to: academic problems, injury, physical/sexual assault, or death


### Studies have found that:

- Nearly one third of college students underestimate the amount of alcohol they plan to drink during spring break
- During spring break vacation at a beach-front destination, the average college male consumes 18 drinks per day, while the average college female consumes 10 drinks per day

To stay safe during spring break, the Centers for Disease Control and Prevention (CDC) recommends that you limit alcohol intake, refrain from drinking and driving, plan ahead, and call 911 if you suspect an overdose.

Wisconsin has a higher percentage of alcohol and binge alcohol use among 18-25 year-olds compared to the U.S.

## Risks & Consequences of Binge Drinking



Source: NSDUH 2016-17

**696,000**  
18-24 year-old U.S. students are assaulted each year by another student who has been drinking

**599,000**  
18-24 year-old U.S. students reported accidental alcohol-related injuries within the past year

**97,000**  
18-24 year-old U.S. students reported being victims of alcohol-related sexual assault within the past year

**1,825**  
18-24 year-old U.S. college students die annually from accidental alcohol-related injuries, including car accidents

Source: SAMHSA, 2017

Sources:  
American College Health Association (ACHA) National College Health Assessment (NCHA) II. (2015). University of Wisconsin System Reference Group Executive Summary Spring 2015.  
Centers for Disease Control and Prevention (CDC). (2015). Spring Break Health and Safety Tips. Retrieved from <https://www.cdc.gov/family/springbreak/index.htm>  
Lee, C., Patrick, M., & Zimmerman, L. (June 2017). Individual, Interpersonal, & Contextual Factors Associated with Discrepancies between Intended and Actual Spring Break Drinking. *Journal of Addictive Behaviors*, 69, 42-47.  
National Institute on Alcohol Abuse and Alcoholism (NIAAA). Drinking Levels Defined. Retrieved from <https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/moderate-binge-drinking>  
National Survey on Drug Use and Health (NSDUH). Substance Abuse and Mental Health Services Administration (SAMHSA), U.S. Department of Health and Human Services (HHS). (2013-14, 2016-17).  
Smeaton, G., Josiam, B., & Dietrich, U. (1998). College Students' Binge Drinking at a Beach-Front Destination During Spring Break. *Journal of American College Health*, 46(6), 247-254.  
SAMHSA. (September 20, 2017). Underage Drinking and School and Campus Health. Retrieved from <https://www.samhsa.gov/underage-drinking-topic/school-campus-health>  
SAMHSA. (August 16, 2016). The CBHSQ Report- Trends in Perception of Risk and Availability of Substance Use Among Full-Time College Students. Retrieved from [https://www.samhsa.gov/data/sites/default/files/report\\_2418/ShortReport-2418.html](https://www.samhsa.gov/data/sites/default/files/report_2418/ShortReport-2418.html)